

Facial Palsy due to Parotid Abscess: An Unusual Complication

Parotis Apsesine Bağlı Yüz Felci: Alışılmadık bir Komplikasyon

Case Report
 Olgu Sunumu

Mehtab Alam, Syed Abrar Hasan, Shahab Farkhund Hashmi, Piyush Kant Singh
 Department of Otorhinolaryngology, Aligarh Muslim University School of Medicine, Aligarh, India

Abstract

Facial nerve palsy is usually associated with a malignant parotid neoplasm; it is highly unusual for it to result from a benign situation, such as inflammation or infection of the parotid gland. Surgery along with prompt medical treatment is the mainstay, and in the majority of the patients, nerve paralysis recovers in the follow-up period. We report a case of a 50-year-old non-diabetic non-hy-

pertensive female who presented with odynophagia, left-sided parotid swelling, and left facial nerve palsy. The diagnosis of facial nerve palsy due to a parotid abscess extending to the parapharyngeal space was made. Facial palsy fully recovered within 2 months.

Keywords: Facial palsy, parotid abscess, parapharyngeal extension

Öz

Fasiyal sinir felci, sıklıkla parotis malign neoplazmları ile ilişkili bir durumdur. Glandda, enflamasyon veya enfeksiyon gibi benign durumlardan kaynaklanan fasiyal sinir felci nadirdir. Acil medikal tedavi ile birlikte cerrahi, asıl tedaviyi oluşturur ve bu hastaların çoğunun izlemine sinir paralizisi iyilesir. Burada, diabet ve hipertansiyonu olmayan, odinofaji, sol parotis bezinde şişlik

ve fasiyal sinir felci ile başvuran 50 yaşında kadın hasta sunuldu. Parafarengeal bölgeye uzanan parotis apsesi nedeniyle gelişen fasiyal sinir felci tanısı konuldu. Yüz felcinin iki ay içinde tamamen iyileştiği gözlemlendi.

Anahtar Kelimeler: Yüz felci, parotis apsesi, parafarengeal uzanım

Introduction

Facial nerve paralysis due to parotid pathology is almost always caused by malignant neoplasm of the gland. Though the benign conditions of the parotid gland, such as benign mixed tumors, Warthin's tumor, sarcoidosis, and parotid cysts can affect the facial nerve, occurrence of facial nerve palsy as a result of a parotid abscess is exceedingly rare, with only nine previously reported cases (1). Parotid abscesses may arise from ductal ectasia, primary parenchymal involvement, or infection of the intra-parotid lymph nodes. Although a conservative treatment with hydration and administration of broad-spectrum antibiotics are the first choice, surgical intervention may be required in patients with no response to medical treatment (2).

Case Report

A 50-year-old non-diabetic non-hypertensive female was admitted to the emergency department of our hospital with complaints of diffuse swelling in the left parotid region and pain during mastication and swallowing for the past 5 days. Symptoms of left facial nerve palsy started just 3 hours before admission, which forced her to seek medical help; she had not taken any medication till then. There was no history of dental pain or ear discharge.

On examination, she was febrile and the general appearance of the patient was not good. There was a diffuse, non-fluctuant, tense, and tender swelling of about 6-7 cm in the left parotid region. The ear lobule was pushed forward. There was facial nerve paralysis of grade 4 (House-Brackmann classification), as the patient was having



Address for Correspondence/Yazışma Adresi:

Mehtab Alam
 E-mail: mehtabalam4105@gmail.com

Received Date/Geliş Tarihi: 11.04.2016

Accepted Date/Kabul Tarihi: 28.09.2016

© Copyright 2016 by Official Journal of the Turkish Society of Otorhinolaryngology and Head and Neck Surgery Available online at
www.turkarchotorhinolaryngol.org

© Telif Hakkı 2016 Türk Kulak Burun Boğaz ve Baş Boyun Cerrahisi Derneği Makale metnine
www.turkarchotorhinolaryngol.org web sayfasından ulaşılabilir.

DOI: 10.5152/tao.2016.1671

incomplete eye closure, asymmetry of the mouth, and no motion in the forehead (Figure 1). Restricted mouth opening due to trismus was present, and examination of oral cavity showed medial bulge of the left lateral pharyngeal wall, displacing the uvula and tonsillar pillar (Figure 2). Complete blood count revealed leukocytosis (16500/ μ L.) with predominant neutrophils (91%).

An urgent contrast-enhanced computed tomography (CT) scan of the parotid region was performed, which was suggestive of a left parotid abscess with parapharyngeal extension (Figure 3, 4). Broad-spectrum intravenous antibiotics (IV co-amoxiclav 1.2 g three times over 24 hours, amikacin 500 mg twice daily, and metronidazole 500 mg/100 mL three time daily) were started immediately along with intravenous rehydration.

Incision and drainage of the abscess was performed under general anesthesia, evacuating around 80 cc of pus, the culture of which showed mixed growth comprising of *Fusobacterium*, *Bacteroides fragilis*, *Staphylococcus aureus*, and *Klebsiella*. Antibiogram results showed *Fusobacterium* and *Bacteroides fragilis* sensitive to metronidazole and amoxicillin, and both *Staphylococcus aureus* and *Klebsiella* were found to be sensitive to amikacin. In the immediate post-operative period, the patient's symptoms were partially relieved, except for facial palsy. Daily dressing was done for 10 days, and on the 11th day, she was discharged with closure of the wound, which was done using non-absorbable sutures; she had slight improvement in the facial paresis at the time of discharge (Figure 5). The patient was called for regular follow-up visits, and within 2 months, her facial nerve palsy showed complete recovery.

Written informed consent was taken from the patient who participated in this study.



Figure 2. Parapharyngeal extension of parotid abscess as evident by the medial push of the uvula and tonsillar pillar



Figure 3. CT image showing left-sided parotid abscess



Figure 1. Left-sided parotid swelling with signs of facial nerve palsy




Figure 4. CT image showing left-sided parotid abscess



Figure 5. Clinical image on the 11th post-operative (discharge) day showing slight improvement in facial paresis

Discussion

Parotid abscess usually occurs in elderly immunocompromised persons who are debilitated with systemic illness (3). Tan and Goh (4), in a series of 15 patients with parotid abscess, revealed that the mean age of presentation was 51 years and that six of them were diabetic. Even in neonates and infants, suppurative parotitis with facial nerve palsy has been described (2, 5). The predisposing factors can be poor oral hygiene, dehydration, and obstruction of Stenson's duct. The most likely route for the entry of the causative organism is ascending migration from the oral cavity to the salivary duct.

It is extremely rare to have facial nerve palsy due to a parotid abscess. In total, there are around 16 reported cases of involvement of the facial nerve in benign and inflammatory conditions of the parotid gland, and only nine of them are due to a parotid abscess (1, 6). The nerve is either partially or completely affected resulting in varied degrees of paralysis.

Most of the parotid abscesses resulting in facial palsy are because of bacteria, with *S. aureus* being the commonest, but there is a report mentioning *Candida albicans* parotid abscess with facial nerve dysfunction (7). Even viral parotitis (mumps) can cause facial palsy (8).

The exact mechanism for involvement of the facial nerve in non-malignant conditions of the parotid gland, as in this case, remains unclear. It had been postulated that the nerve is affected due to perineuritis and local toxic effects along with ischemic neuropathy because of rapidly increasing swelling causing pressure over the nerve (9).

The CT scan is the imaging modality of choice, as there may be difficulty in differentiating between acute parotitis and parotid abscess because of the tense parotid fascia. Sometimes, an underlying malignancy is detected on CT, which causes facial nerve palsy (10). Contrast-enhanced CT in our patient showed parotid abscess with parapharyngeal extension.

In the pre-antibiotic era, parotid abscesses were known for poor outcomes with a high mortality, which has dramatically come down nowadays due to use of proper antibiotics (11). Once the diagnosis of parotid abscess is established, surgical treatment (incision and drainage) along with broad-spectrum antibiotics should always be taken up and is the treatment of choice (1).

The Blair technique is used for incision and drainage (12). A pre-auricular retromandibular incision is given; flaps of skin and subcutaneous tissue are reflected superficial to the parotid fascia; and then, using a hemostat, multiple openings are made always parallel to the branches of the facial nerve.

Facial nerve palsy in our patient recovered fully within 2 months, although there have been cases with residual paralysis even after 6 months (3).

Conclusion

Facial nerve palsy as a complication of parotid abscess is very rare. Radiological imaging, such as the CT scan, greatly helps in making an early diagnosis after which the treatment is initiated, with surgery being the mainstay along with broad-spectrum antibiotics. In most of the cases, there is complete recovery of facial nerve dysfunction.

Informed Consent: Written informed consent was obtained from patient who participated in this study.

Authors contribution: Concept - M.A., S.A.H.; Design - M.A., P.K.S.; Supervision - S.A.H., S.F.H.; Resource - M.A., S.F.H.; Material - M.A., P.K.S.; Data Collection and/or Processing - M.A., S.A.H., P.K.S.; Analysis and/or Interpretation - M.A., S.F.H., P.K.S.; Literature Search - M.A., S.A.H., P.K.S.; Writing - M.A., P.K.S.; Critical Reviews - S.A.H., S.F.H.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

Hasta Onamı: Yazılı hasta onamı bu çalışmaya katılan hastadan alınmıştır.

Hakem Değerlendirmesi: Dış bağımsız.

Yazar Katkıları: Fikir - M.A., S.A.H.; Tasarım - M.A., P.K.S.; Denetleme - S.A.H., S.F.H.; Kaynaklar - M.A., S.F.H.; Gereçler - M.A., P.K.S.; Veri Toplanması ve/veya İşlemesi - M.A., S.A.H., P.K.S.; Analiz ve/veya Yorum - M.A., S.F.H., P.K.S.; Literatür Taraması - M.A., S.A.H., P.K.S.; Yazıyı Yazan - M.A., P.K.S.; Eleştirel İnceleme - S.A.H., S.F.H.

Çıkar Çatışması: Yazarlar çıkar çatışması bildirmemişlerdir.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

References

1. Kristensen RN, Hahn CH. Facial nerve palsy caused by parotid abscess. *J Laryngol Otol* 2012; 126: 322-4. [\[CrossRef\]](#)
2. Anitha C, Kumar KJ, Kumar MG, Chavan A. Facial nerve palsy caused by parotid abscess in an infant. *Indian J Pediatr* 2014; 81: 1251-2. [\[CrossRef\]](#)
3. Sabir Husin Athar PP, Yahya Z, Mat Baki M, Abdullah A. Facial nerve paralysis: a rare complication of parotid abscess. *Malays J Med Sci* 2009; 16: 39-40.
4. Tan VE, Goh BS. Parotid abscess: a five year review- clinical presentation, diagnosis and management. *J Laryngol Otol* 2007; 12: 872-9. [\[CrossRef\]](#)
5. Mathur NB, Goyal RK, Khalil A. Neonatal suppurative parotitis with facial palsy. *Indian Pediatrics* 1988; 25: 806-7.
6. Orhan KS, Demirel T, Kocasoy-Orhan E, Yenigül K. Facial paralysis due to occult parotid abscess. *Kulak Burun Bogaz Ihtis Derg* 2008; 18: 115-7.
7. Marioni G, Rinaldi R, de Filippis C, Gaio E, Staffieri A. Candidal abscess of parotid gland associated with facial nerve paralysis. *Acta Otolaryngol* 2003; 123: 661-3. [\[CrossRef\]](#)
8. Incecik F, Herguner MO, Altunbasak S. Facial palsy caused by mumps parotitis. *Neurol India* 2009; 57: 511-2. [\[CrossRef\]](#)
9. Smith DR, Hartig GK. Complete facial paralysis as a result of parotid abscess. *Otolaryngol Head Neck Surg* 1997; 117: S114-7. [\[CrossRef\]](#)
10. Casselman JW, Mancuso AA. Major salivary gland masses: Comparison of MR imaging and CT. *Radiology* 1987; 165: 183-9. [\[CrossRef\]](#)
11. Raad I, Sabbagh MF, Caranasos GJ. Acute bacterial sialadenitis: A study of 29 cases and review. *Reviews of Infectious Diseases* 1990; 12: 591-601. [\[CrossRef\]](#)
12. Blair VP, Pagett EC. Pyogenic infection of the parotid glands and ducts. *Arch Surg* 1923; 7: 1-36. [\[CrossRef\]](#)